

## Comment on Liu et al.: Operative versus non-operative treatment for clavicle fracture: a meta-analysis

Chenli Ye · Guoping Liao · Shuai He · Zhongyi Zhang

Received: 12 May 2013 / Accepted: 4 June 2013 / Published online: 22 June 2013  
© The Author(s) 2013. This article is published with open access at Springerlink.com

Dear Editor,

We read with great interest the recent article entitled “Operative versus non-operative treatment for clavicle fracture: a meta-analysis” published online in May 2013 issue of International Orthopaedics by Liu et al. [1]. The authors performed a meta-analysis to evaluate operative versus non-operative treatment for clavicle fractures. They reached an important conclusion that operative treatment is better than non-operative treatment. It is a valuable study. Nevertheless, there are some comments we would like to raise related to this article.

1. The authors clearly stated that “Five studies involving 633 clavicle fractures were included” in the abstract. However, we found eight studies were included in the subsequent pooled analysis. And, it is not in accordance with the original study by Smekal et al. [2] that 68 patients were included in this article by Liu et al. [1], as shown in Table 1. Actually, the number of patients reported by Smekal et al. [2] was 60. Therefore, we conclude that five RCTs and three CCTs involving 655 clavicle fractures should be included.
2. Four electronic databases (MEDLINE, EMBASE, OVID and the Cochrane Central Register of Controlled Trials) were systematically searched by the authors. However, using the same search strategy and end of search date as those of Liu et al., we could not find two relevant studies [3, 4] that have been included in the meta-analysis even though they satisfied the search criteria.

Therefore, the authors should focus specifically or in detail on the issue of the completeness of the search strategy report for databases. The search strategy report plays an important role in systematic reviews.

3. The authors have clearly stated that “The quality of the included studies was assessed by two authors” in the abstract. We consider that the authors could evaluate methodological quality for all selected studies (randomised controlled trials and controlled clinical trials), which could avoid the potential bias in this meta-analysis. However, the authors did not describe how to assess the quality of all studies in the meta-analysis and there were no detailed scores for each trial.
4. There are different approaches in either operative or nonoperative treatment for clavicle fracture. If possible, we suggest that a meta-analysis of different operative approaches versus different nonoperative treatment for clavicle fracture could be conducted.
5. The authors have clearly stated that “The difference in neurological complications (RR 0.45, 95%CI 0.25–0.81) was statistically significant between operative and non-operative treatment” in the abstract, and then they write “there was also no statistical difference in the neurological complication rate between operative and non-operative treatment” in the results part. We suggest that “no statistical difference” should be replaced by “statistical difference” in the results section. Meanwhile, the result of the forest plots for neurological complications showed that the difference was statistically significant between operative and non-operative treatment.
6. Although both techniques have been associated with excellent results in a variety of reviews, the lack of a standard operative protocol must be considered a weakness of the meta-analysis; it may be that there are fundamental differences in outcome between the two techniques. In addition, patients with a complete clavicular fracture

C. Ye · S. He · Z. Zhang (✉)  
Department of Pharmacy, Zhujiang Hospital, Southern Medical University, Guangzhou 510282, People's Republic of China  
e-mail: zhang43499@sohu.com

G. Liao  
Changning Hospital of Traditional Chinese Medicine, Changning 421500, Hunan Province, People's Republic of China

may be counselled that they will be at a higher risk of sustaining nonunion and symptomatic malunion if the fracture is treated nonoperatively, since there is no clear evidence that surgical treatment will improve their long-term function in general.

In conclusion, the results of this meta-analysis by Liu et al. [1] should be interpreted with caution. To reach a definitive conclusion, further high quality RCTs based on larger sample sizes are still needed to assess the effects of operative and non-operative treatment on clavicle fractures. We believe that our remarks will contribute to a more accurate elaboration of the results presented by Liu et al. (2013).

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

1. Liu GD, Tong SL, Ou S, Zhou LS, Fei J, Nan GX, Gu JW (2013) Operative versus non-operative treatment for clavicle fracture: a meta-analysis. *Int Orthop*. doi:10.1007/s00264-013-1871-z
2. Smekal V, Irenberger A, Struve P, Wambacher M, Krappinger D, Kralinger FS (2009) Elastic stable intramedullary nailing versus nonoperative treatment of displaced midshaft clavicular fractures-a randomized, controlled, clinical trial. *J Orthop Trauma* 23(2):106–112
3. Smith C, Rudd J, Crosby L (2001) Results of operative versus non-operative treatment for 100% displaced mid-shaft clavicle fractures: a prospective randomized trial. In: *Proceedings from the 68th Annual Meeting of the American Academy of Orthopaedic Surgeons*, San Francisco, CA
4. Virtanen KJ, Paavola MP, Remes VM, Pajarinen J, Savolainen V, Bjorkenheim JM (2010) Nonoperative versus operative treatment of midshaft clavicle fractures: a randomized controlled trial. *The 75th Annual Meeting of the American Academy of Orthopaedic Surgeons*, San Francisco, CA, 9–12, pp 65–73